

# Zakkaart - Sedatie algoritme voor patiënten met acute gedragsstoornissen op de SEH

## Referenties:

1. Richtlijn Intoxicaties: eerste opvang in het ziekenhuis. 2017. <https://internisten.nl/sites/internisten.nl/files/Richtlijn-intoxicaties-18122017-geautoriseerd.pdf>
2. Management of patients with Acute Severe Behavioural Disturbance in Emergency Departments - Guideline. NSW Health. 2015. (Review date 10-Aug-2020)
3. Isbister GK, Calver LA, Page CB, Stokes B, Bryant JL, Downes MA. Randomized controlled trial of intramuscular droperidol versus midazolam for violence and acute behavioral disturbance: the DORM study. Ann Emerg Med. 2010 Oct;56(4):392-401.e1.
4. Calver L, Page CB, Downes MA, Chan B, Kinnear F, Wheatley L, Spain D, Isbister GK. The Safety and Effectiveness of Droperidol for Sedation of Acute Behavioral Disturbance in the Emergency Department. Ann Emerg Med. 2015 Sep;66(3):230-238.e1.
5. Taylor DM, Yap CYL, Knott JC, Taylor SE, Phillips GA, Karro J, Chan EW, Kong DCM, Castle DJ. Midazolam-Droperidol, Droperidol, or Olanzapine for Acute Agitation: A Randomized Clinical Trial. Ann Emerg Med. 2017 Mar;69(3):318-326.e1.
6. Chan EW, Taylor DM, Knott JC, Phillips GA, Castle DJ, Kong DC. Intravenous droperidol or olanzapine as an adjunct to midazolam for the acutely agitated patient: a multicenter, randomized, double-blind, placebo-controlled clinical trial. Ann Emerg Med. 2013 Jan;61(1):72-81.
7. Cole JB, Lee SC, Martel ML, Smith SW, Biros MH, Miner JR. The Incidence of QT Prolongation and Torsades des Pointes in Patients Receiving Droperidol in an Urban Emergency Department. West J Emerg Med. 2020 Jul 2;21(4):728-736.
8. Kim HK, Leonard JB, Corwell BN, Connors NJ. Safety and efficacy of pharmacologic agents used for rapid tranquilization of emergency department patients with acute agitation or excited delirium. Expert Opin Drug Saf. 2021 Feb;20(2):123-138.